

Heat Kernel for Open Manifolds

Trevor H. Jones

October 23, 2009

Abstract

In a 1991 paper by Buttig and Eichhorn, the existence and uniqueness of a differential forms heat kernel on open manifolds of bounded geometry was proven. In that paper, it was shown that the heat kernel obeyed certain properties, one of which was a relationship between the derivative of heat kernel of different degrees. We will give a proof of this condition for complete manifolds with Ricci curvature bounded below, and then use it to give an integral representation of the heat kernel of degree k .

In this paper we are considering the differential forms heat equation on manifolds, in particular we are considering $(\Delta + \partial_t)\omega = 0$ with Dirichlet initial conditions. Our goal is to produce a formula for the Green's function, also known as the heat kernel or fundamental solution, which gives the solution of this equation.

The solutions of this equation in the case of functions, or 0-forms, is well-known. The work on differential forms has been much more recent. In 1983, Dodziuk [Dod83] proved that for complete oriented C^∞ Riemannian manifolds with Ricci curvature bounded below, bounded solutions are uniquely determined by their initial values. In 1991 Buttig and Eichhorn [BE91] gave existence and uniqueness results for the differential forms heat kernel on open manifolds of bounded geometry. In the same paper they also gave an identity relating derivatives of the heat kernel of differing degrees. Using that identity, the author, in his doctoral thesis, [Jon08], was able to give a formula for the 1-form heat kernel on open Riemann surfaces of bounded geometry. This formula,

$$K_1(\mathbf{x}, \mathbf{y}, t) = (I + *_{\mathbf{x}} *_{\mathbf{y}}) d_{\mathbf{x}} d_{\mathbf{y}} \int_t^\infty K_0(\mathbf{x}, \mathbf{y}, \tau) d\tau, \quad (1)$$

directly relates the 1-form heat kernel to the 0-form heat kernel, about which more is known.

One of the properties given by Buttig and Eichhorn for a global heat kernel was that the heat kernels $K_k(\mathbf{x}, \mathbf{y}, t)$ and $K_{k+1}(\mathbf{x}, \mathbf{y}, t)$ are related by

$$d_{\mathbf{x}}K_k(\mathbf{x}, \mathbf{y}, t) = d_{\mathbf{y}}^*K_{k+1}(\mathbf{x}, \mathbf{y}, t). \quad (2)$$

In this article, we will present an alternate proof of this property for manifolds with Ricci curvature bounded below, and then use this to give a formula for the k -form heat kernel.

Lemma 1 *For a complete manifold M , with Ricci curvature bounded from below, we have the following relationship between the k - and $(k+1)$ -form heat kernels:*

1. $d_{\mathbf{x}}K_k(\mathbf{x}, \mathbf{y}, t) = d_{\mathbf{y}}^*K_{k+1}(\mathbf{x}, \mathbf{y}, t)$
2. $d_{\mathbf{y}}K_k(\mathbf{x}, \mathbf{y}, t) = d_{\mathbf{x}}^*K_{k+1}(\mathbf{x}, \mathbf{y}, t)$

Proof: Let $E(\mathbf{x}, \mathbf{y}, t) = d_{\mathbf{x}}K_k(\mathbf{x}, \mathbf{y}, t) - d_{\mathbf{y}}^*K_{k+1}(\mathbf{x}, \mathbf{y}, t)$. We will demonstrate that E satisfies the heat equation with zero as the initial condition. This will imply, by uniqueness of the solutions of the heat equation, see [Dod83], that $E \equiv 0$, giving the desired result.

First

$$\begin{aligned} \Delta_{\mathbf{x}}E &= \Delta_{\mathbf{x}}d_{\mathbf{x}}K_k(\mathbf{x}, \mathbf{y}, t) - \Delta_{\mathbf{x}}d_{\mathbf{y}}^*K_{k+1}(\mathbf{x}, \mathbf{y}, t) \\ &= d_{\mathbf{x}}\Delta_{\mathbf{x}}K_k(\mathbf{x}, \mathbf{y}, t) - d_{\mathbf{y}}^*\Delta_{\mathbf{x}}K_{k+1}(\mathbf{x}, \mathbf{y}, t) \\ &= d_{\mathbf{x}}(-\partial_t)K_k(\mathbf{x}, \mathbf{y}, t) - d_{\mathbf{y}}^*(-\partial_t)K_{k+1}(\mathbf{x}, \mathbf{y}, t) \\ &= -\partial_t E. \end{aligned}$$

Next consider $W := \langle E, \omega(\mathbf{x}) \rangle$, where ω is a suitable test function and $\langle \mu, \nu \rangle = \int_M \mu \wedge * \nu$. Then

$$\begin{aligned} \lim_{t \rightarrow 0+} W &= \lim_{t \rightarrow 0+} \langle d_{\mathbf{x}}K_k, \omega(\mathbf{x}) \rangle - \langle d_{\mathbf{y}}^*K_{k+1}, \omega(\mathbf{x}) \rangle \\ &= \lim_{t \rightarrow 0+} \langle K_k, d_{\mathbf{x}}^*\omega(\mathbf{x}) \rangle - d_{\mathbf{y}}^* \langle K_{k+1}, \omega(\mathbf{x}) \rangle \\ &= d_{\mathbf{y}}^*\omega(\mathbf{y}) - d_{\mathbf{y}}^*\omega(\mathbf{y}) = 0 \end{aligned}$$

Since ω was an arbitrary test function, we must have that $E \equiv 0$ at $t = 0$. Thus by uniqueness, $E \equiv 0$ for all $t > 0$.

The proof of the second assertion follows in a similar manner. ■

We will use this result to give an explicit formula for K_k in terms of $K_{k\pm 1}$.

Theorem 2 *Let M be an open, complete manifold with Ricci curvature bounded below. Then the differential forms heat kernel obey the following relation:*

$$K_k(\mathbf{x}, \mathbf{y}, t) = d_{\mathbf{x}} d_{\mathbf{y}} \int_t^\infty K_{k-1}(\mathbf{x}, \mathbf{y}, \tau) d\tau + d_{\mathbf{x}}^* d_{\mathbf{y}}^* \int_t^\infty K_{k+1}(\mathbf{x}, \mathbf{y}, \tau) d\tau.$$

Proof: Let K_k be the k -form heat kernel. Clearly,

$$K_k(\mathbf{x}, \mathbf{y}, t) = - \int_t^\infty \frac{\partial}{\partial \tau} K_k(\mathbf{x}, \mathbf{y}, \tau) d\tau,$$

since K_k tends to zero (pointwise) as t increases. Since K_k is a solution of the heat equation, we can replace the time derivative with $-\Delta_{\mathbf{x}} = -d_{\mathbf{x}} d_{\mathbf{x}}^* - d_{\mathbf{x}}^* d_{\mathbf{x}}$, so

$$K_k(\mathbf{x}, \mathbf{y}, t) = \int_t^\infty (d_{\mathbf{x}} d_{\mathbf{x}}^* + d_{\mathbf{x}}^* d_{\mathbf{x}}) K_k(\mathbf{x}, \mathbf{y}, \tau) d\tau.$$

Using Lemma 1, we can rewrite the above as

$$K_k(\mathbf{x}, \mathbf{y}, t) = \int_t^\infty d_{\mathbf{x}} d_{\mathbf{y}} K_{k-1}(\mathbf{x}, \mathbf{y}, \tau) + d_{\mathbf{x}}^* d_{\mathbf{y}}^* K_{k+1}(\mathbf{x}, \mathbf{y}, \tau) d\tau. ■$$

The result in Theorem 2 depends mainly on two things: the existence and uniqueness of the heat kernel, and the pointwise convergence to zero of the kernel for large time. The methods used above work for a diffusion-type equation provided these conditions are met. For example, for the diffusion equation $(\Delta + c\partial_t)\omega = 0$, the proofs follow through almost identically.

Corollary 3 *Let M be an open, n -dimensional, differentiable manifold, with Ricci curvature bounded below, and consider the differential form diffusion equation $(\Delta + c\partial_t)\omega = 0$ with initial data $\omega(\mathbf{x}, 0) = f(\mathbf{x})$. Then the Green's functions are related by*

$$G_k(\mathbf{x}, \mathbf{y}, t) = d_{\mathbf{x}} d_{\mathbf{y}} \int_{ct}^\infty G_{k-1}(\mathbf{x}, \mathbf{y}, \tau) d\tau + d_{\mathbf{x}}^* d_{\mathbf{y}}^* \int_{ct}^\infty G_{k+1}(\mathbf{x}, \mathbf{y}, \tau) d\tau.$$

Proof: Let $T = ct$, then $c\partial_t = \partial_T$, so the equation becomes $(\Delta + \partial_T)\omega(\mathbf{x}, T) = 0$ with the same initial conditions. So by Theorem 2 we have the desired Green's functions. $\blacksquare$

In the case of 2-dimensional manifolds, the 0-form and the 2-form heat kernels are isomorphic, as the following Lemma will show. This allows use to write the 1-form heat kernel in terms of the 0-form, or function, heat kernel.

Lemma 4 *Let M be a complete manifold with Ricci curvature bounded below. Then the differential forms heat kernels, K_k and K_{n-k} are related in the following manner:*

$$K_k = *_x *_y K_{n-k}.$$

Proof: Consider the equation $(\partial_t + \Delta_k)u = 0$, $u(\mathbf{x}, 0) = f(\mathbf{x})$. Then u is given by $u(\mathbf{x}, t) = \langle K_k(\mathbf{x}, \mathbf{y}, t), f(\mathbf{y}) \rangle = \int_M K(\mathbf{x}, \mathbf{y}, t) \wedge_{\mathbf{y}} *_y f(\mathbf{y})$. Since $*\Delta_k = \Delta_{n-k}*$, it follows that $*_{\mathbf{x}}u$ is a solution of the $(n-k)$ -form heat equation with initial condition $*_{\mathbf{x}}f(\mathbf{x})$. So

$$\begin{aligned} *_x u(\mathbf{x}, t) &= \langle K_{n-k}(\mathbf{x}, \mathbf{y}, t), *_y f(\mathbf{y}) \rangle \\ &= \langle *_y f(\mathbf{y}), K_{n-k}(\mathbf{x}, \mathbf{y}, t) \rangle \\ &= \int_M *_y f(\mathbf{y}) \wedge_{\mathbf{y}} *_y K_{n-k}(\mathbf{x}, \mathbf{y}, t) \end{aligned}$$

By applying $*_{\mathbf{x}}$ to both sides, and changing order in the wedge product, we have

$$(-1)^{k(n-k)}u(\mathbf{x}, t) = \int_M (-1)^{k(n-k)} *_x *_y K_{n-k}(\mathbf{x}, \mathbf{y}, t) \wedge_{\mathbf{y}} *_y f(\mathbf{y})$$

or

$$u(\mathbf{x}, t) = \langle *_x *_y K_{n-k}(\mathbf{x}, \mathbf{y}, t), f(\mathbf{y}) \rangle.$$

By uniqueness of the heat kernel we have the desired result. $\blacksquare$

Corollary 5 *Let M be an open, complete manifold of dimension 2 with Ricci curvature bounded below. Then the 1-form heat kernel on M is given by*

$$K_1(\mathbf{x}, \mathbf{y}, t) = (I + *_x *_y) d_{\mathbf{x}} d_{\mathbf{y}} \int_t^\infty K_0(\mathbf{x}, \mathbf{y}, \tau) d\tau$$

where, $\mathbf{x}, \mathbf{y} \in M$ and $t > 0$ and K_0 is the 0-form heat kernel.

Proof: Since M has dimension 2, and so $K_2 = *_x *_y K_0$ by Lemma 4. Recall that $d^* * \omega = - * d\omega$ for 0-forms, ω . This gives the desired result. $\blacksquare$

As an example, consider the case of the hyperbolic plane, with constant curvature -1 . From [Cha84] we have the 0-form heat kernel

$$K_0(\mathbf{x}, \mathbf{y}, t) = \frac{1}{2\pi} \int_0^\infty P_{-\frac{1}{2}+i\rho}(\cosh d_{H^2}(\mathbf{x}, \mathbf{y})) \rho e^{-(\frac{1}{4}+\rho^2)t} \tanh \pi \rho d\rho,$$

which, if we perform the integration set out in Corollary 5, we get

$$K_1(\mathbf{x}, \mathbf{y}, t) = \frac{1}{2\pi} (I + *_x *_y) d_x d_y \left[\int_0^\infty P_{-\frac{1}{2}+i\rho}(\cosh d_{H^2}(\mathbf{x}, \mathbf{y})) \rho \frac{e^{-(\frac{1}{4}+\rho^2)t}}{\frac{1}{4}+\rho^2} \tanh \pi \rho d\rho \right].$$

If M is an open 2-dimensional manifold which has a unique heat kernel for functions, K_0 , then Corollary 5 suggests a candidate for a heat kernel on 1-forms, and since the K_0 and K_2 heat kernels are isomorphic, we would know all the heat kernels. We will now show that $K_1(\mathbf{x}, \mathbf{y}, t) = (I + *_x *_y) d_x d_y \int_t^\infty K_0(\mathbf{x}, \mathbf{y}, \tau) d\tau$ works as the heat kernel. So given

$$(\Delta_{\mathbf{x}}^{(1)} + \partial_t) w_1(\mathbf{x}, t) = 0 \tag{3}$$

$$w_1(\mathbf{x}, 0) = f_1(\mathbf{x}) \tag{4}$$

show that w_1 can be written as $w_1(\mathbf{x}, t) = \langle K_1(\mathbf{x}, \mathbf{y}, t), f_1(\mathbf{y}) \rangle$.

So, let w_1 be a solution of (3) and (4), and $w(\mathbf{x}, t) = \langle K_1(\mathbf{x}, \mathbf{y}, t), f_1(\mathbf{y}) \rangle$. Since the Laplacian commutes with the Hodge star isomorphism and the exterior derivative and coderivative, it is clear that w satisfies equation (3). Now we just need to show that w as defined, satisfies the initial condition (4).

$$\begin{aligned} w(\mathbf{x}, t) &= \int_t^\infty d_x \langle d_y K_0(\mathbf{x}, \mathbf{y}, \tau), f_1(\mathbf{y}) \rangle + *_x d_x \langle *_y d_y K_0(\mathbf{x}, \mathbf{y}, \tau), f_1(\mathbf{y}) \rangle d\tau \\ &= \int_t^\infty d_x \langle K_0(\mathbf{x}, \mathbf{y}, \tau), d_y^* f_1(\mathbf{y}) \rangle - *_x d_x \langle K_0(\mathbf{x}, \mathbf{y}, \tau), *_y d_y f_1(\mathbf{y}) \rangle d\tau \\ &= \int_t^\infty d_x d_x^* w_1(\mathbf{x}, \tau) - *_x d_x *_x d_x w_1(\mathbf{x}, \tau) d\tau \\ &= \int_t^\infty \Delta w_1(\mathbf{x}, \tau) d\tau \\ &= \int_t^\infty -\partial_\tau w_1(\mathbf{x}, \tau) d\tau = w_1(\mathbf{x}, t) \end{aligned}$$

Since w_1 is a solution of the heat equation with initial value f_1 , and $w = w_1$, this means that w also has initial value f_1 . Thus w is a solution of (3) and (4).

Finally, let us consider the case of compact complete manifolds. In this case, because of conservation, diffusion does not tend to zero, so the large-time limit has to be taken into account.

Theorem 6 *Let M be a complete manifold with Ricci curvature bounded below, and that the $\lim_{t \rightarrow \infty} K_k(\mathbf{x}, \mathbf{y}, t)$ is a constant double-form, call it C . Then, the heat kernel obeys the following relation:*

$$K_k(\mathbf{x}, \mathbf{y}, t) = C + d_{\mathbf{x}} d_{\mathbf{y}} \int_t^\infty K_{k-1}(\mathbf{x}, \mathbf{y}, \tau) d\tau + d_{\mathbf{x}}^* d_{\mathbf{y}}^* \int_t^\infty K_{k+1}(\mathbf{x}, \mathbf{y}, \tau) d\tau.$$

Proof: Let K_k be the k -form heat kernel. Clearly,

$$K_k(\mathbf{x}, \mathbf{y}, t) = C - \int_t^\infty \frac{\partial}{\partial \tau} K_k(\mathbf{x}, \mathbf{y}, \tau) d\tau,$$

since K_k tends to C as t increases. Since K_k is a solution of the heat equation, we can replace the time derivative with $-\Delta_{\mathbf{x}} = -d_{\mathbf{x}} d_{\mathbf{x}}^* - d_{\mathbf{x}}^* d_{\mathbf{x}}$, so

$$K_k(\mathbf{x}, \mathbf{y}, t) = C + \int_t^\infty (d_{\mathbf{x}} d_{\mathbf{x}}^* + d_{\mathbf{x}}^* d_{\mathbf{x}}) K_k(\mathbf{x}, \mathbf{y}, \tau) d\tau.$$

Using Lemma 1, we can rewrite the above as

$$K_k(\mathbf{x}, \mathbf{y}, t) = C + \int_t^\infty d_{\mathbf{x}} d_{\mathbf{y}} K_{k-1}(\mathbf{x}, \mathbf{y}, \tau) + d_{\mathbf{x}}^* d_{\mathbf{y}}^* K_{k+1}(\mathbf{x}, \mathbf{y}, \tau) d\tau.$$

■

I would like to thank Dr. Kucerovsky for his helpful comments of earlier versions of this paper.

References

- [BE91] Ingolf Buttig and Jurgen Eichhorn. The heat kernel for p -forms on manifolds of bounded geometry. *Acta Sci. Math.*, 55:33–51, 1991.

- [Cha84] Isaac Chavel. *Eigenvalues in Riemannian Geometry*, volume 115 of *Pure and Applied Mathematics*. Academic Press, 1984.
- [Dod83] Jozef Dodziuk. Maximum Principle for Parabolic Inequalities and the Heat Flow on Open Manifolds. *Indiana University Mathematics Journal*, 32(5):703–16, 1983.
- [Jon08] Trevor H. Jones. *The Heat Kernel on Noncompact Riemann Surfaces*. PhD thesis, University of New Brunswick, 2008.